
 <p><b>Oregon Health &amp; Science University Hospital and Clinics Provider's Orders</b></p> <div style="display: flex; align-items: center;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg); font-size: small; margin-right: 5px;">PO9031</div>  </div> <p style="text-align: center; font-weight: bold;">ADULT AMBULATORY INFUSION ORDER <b>Golimumab (SIMPONI ARIA) Infusion</b></p> <p style="text-align: center; font-size: small;">Page 1 of 3</p>	<p>ACCOUNT NO. _____</p> <p>MED. REC. NO. _____</p> <p>NAME _____</p> <p>BIRTHDATE _____</p> <p style="text-align: right; font-size: x-small; margin-top: 20px;"><i>Patient Identification</i></p>
<b>ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.</b>	

Weight: \_\_\_\_\_ kg      Height: \_\_\_\_\_ cm

Allergies: \_\_\_\_\_

Diagnosis Code: \_\_\_\_\_

Treatment Start Date: \_\_\_\_\_      Patient to follow up with provider on date: \_\_\_\_\_

**\*\*This plan will expire after 365 days at which time a new order will need to be placed\*\***

**GUIDELINES FOR ORDERING**

1. Send **FACE SHEET and H&P or most recent chart note.**
2. Hepatitis B (Hep B surface antigen and core antibody total) screening must be completed prior to initiation of treatment and the patient should not be infected. Please send results with order.
3. A Tuberculin test must have been placed and read as negative prior to initiation of treatment (PPD or QuantiFERON Gold blood test). Please send results with order. If result is indeterminate, a follow up chest X-ray must be performed to rule out TB. Please send results with order.

**PRE-SCREENING: (Results must be available prior to initiation of therapy):**

- Hepatitis B surface antigen and core antibody total test results scanned with orders.
- Tuberculin skin test or QuantiFERON Gold blood test results scanned with orders.
- Chest X-Ray result scanned with orders if TB test result is indeterminate.

**LABS:**

1. CBC with differential, Routine, ONCE, every visit, If on methotrexate or leflunomide
2. CMP, Routine, ONCE, every visit, If on methotrexate or leflunomide

**NURSING ORDERS:**

1. TREATMENT PARAMETER – Hold treatment and contact provider if Hepatitis B surface antigen or core antibody total test result is positive, TB test result is positive, or if screening has not been performed.
2. VITAL SIGNS – Prior to infusion and at the completion of the infusion.
3. Infusions to be scheduled at weeks 0 and 4, then every 8 weeks thereafter.

**MEDICATIONS: (check all that apply)**

golimumab (SIMPONI ARIA) 2 mg/kg in sodium chloride 0.9%, intravenous, ONCE, over 30 minutes

- Initial doses:** Every 4 weeks for 2 treatments (week 0, 4)
- Maintenance doses:** Every 8 weeks thereafter (week 12 and beyond)

**AS NEEDED MEDICATIONS:**

1. acetaminophen (TYLENOL) tablet, 650 mg, oral, EVERY 4 HOURS AS NEEDED for headache, fever, chills or malaise
2. diphenhydramine (BENADRYL) capsule, 50 mg, oral, EVERY 4 HOURS AS NEEDED for itching



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Golimumab (SIMPONI ARIA)  
Infusion**

Page 2 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.**

**HYPERSENSITIVITY MEDICATIONS:**

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

**By signing below, I represent the following:**

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in:  Oregon  \_\_\_\_\_ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

**My physician license Number is # \_\_\_\_\_ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION);** and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

**Provider signature:** \_\_\_\_\_ **Date/Time:** \_\_\_\_\_

**Printed Name:** \_\_\_\_\_ **Phone:** \_\_\_\_\_ **Fax:** \_\_\_\_\_



Oregon Health & Science University  
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER  
**Golimumab (SIMPONI ARIA)  
Infusion**

Page 3 of 3

ACCOUNT NO.  
MED. REC. NO.  
NAME  
BIRTHDATE

*Patient Identification*

**ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK ( ✓ ) TO BE ACTIVE.**

***Please check the appropriate box for the patient's preferred clinic location:***

**Hillsboro Medical Center**

Infusion Services  
364 SE 8th Ave, Medical Plaza Suite 108B  
Hillsboro, OR 97123  
Phone number: (503) 681-4124  
Fax number: (503) 681-4120

**Adventist Health Portland**

Infusion Services  
10123 SE Market St  
Portland, OR 97216  
Phone number: (503) 261-6631  
Fax number: (503) 261-6756

**Mid-Columbia Medical Center**

Celilo Cancer Center  
1800 E 19th St  
The Dalles, OR 97058  
Phone number: (541) 296-7585  
Fax number: (541) 296-7610