



Oregon Health & Science University
Hospital and Clinics Provider's Orders

PO9031



ADULT AMBULATORY INFUSION ORDER

**Ferric Carboxymaltose
(INJECTAFER) Infusion**

Page 1 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

Weight: _____ kg Height: _____ cm

Allergies: _____

Diagnosis Code: _____

Treatment Start Date: _____ Patient to follow up with provider on date: _____

****This plan will expire after 365 days at which time a new order will need to be placed****

GUIDELINES FOR ORDERING

1. Send **FACE SHEET and H&P or most recent chart note.**
2. **Provider must order and obtain a ferritin prior to patient being scheduled for iron infusion. Labs drawn date: _____**

NURSING ORDERS:

1. TREATMENT PARAMETER – Hold treatment and notify provider if Ferritin greater than 300 ng/mL.
2. Monitor the patient for signs and symptoms of hypersensitivity during the infusion and for at least 30 minutes after completion of the infusion. Also monitor BP following infusion.
3. Instruct patient to set follow up appointment with provider for follow up labs.
4. Follow facility policies and/or protocols for vascular access maintenance with appropriate flush solution, declotting (alteplase), and/or dressing changes.

MEDICATIONS:

ferric carboxymaltose (INJECTAFER): (*must check one*)

- Weight 50 kg or greater – 750 mg in sodium chloride 0.9% 250 mL, intravenous, ONCE, over 30 minutes
- Weight less than 50 kg – 15 mg/kg = _____ mg in sodium chloride 0.9%, intravenous, ONCE, over 30 minutes (Pharmacy to prepare in an appropriate volume)

Avoid extravasation (may cause persistent discoloration). Monitor, if extravasation occurs, discontinue administration at that site.

Interval: (*must check one*)

- 2 doses at least 7 days apart
- Other: _____

AS NEEDED MEDICATIONS:

1. sodium chloride 0.9%, 500 mL, intravenous, AS NEEDED x1 dose for vein discomfort. Give concurrently with ferric carboxymaltose



Oregon Health & Science University
Hospital and Clinics Provider's Orders

ADULT AMBULATORY INFUSION ORDER

**Ferric Carboxymaltose
(INJECTAFER) Infusion**

Page 2 of 2

ACCOUNT NO.
MED. REC. NO.
NAME
BIRTHDATE

Patient Identification

ALL ORDERS MUST BE MARKED IN INK WITH A CHECKMARK (✓) TO BE ACTIVE.

HYPERSENSITIVITY MEDICATIONS:

1. NURSING COMMUNICATION – If hypersensitivity or infusion reactions develop, temporarily hold the infusion and notify provider immediately. Administer emergency medications per the Treatment Algorithm for Acute Infusion Reaction (OHSU HC-PAT-133-GUD, HMC C-132). Refer to algorithm for symptom monitoring and continuously assess as grade of severity may progress.
2. diphenhydrAMINE (BENADRYL) injection, 25-50 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
3. EPINEPHrine HCl (ADRENALIN) injection, 0.3 mg, intramuscular, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
4. hydrocortisone sodium succinate (SOLU-CORTEF) injection, 100 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction
5. famotidine (PEPCID) injection, 20 mg, intravenous, AS NEEDED x 1 dose for hypersensitivity or infusion reaction

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*);

I hold an active, unrestricted license to practice medicine in: Oregon _____ (*check box that corresponds with state where you provide care to patient and where you are currently licensed. Specify state if not Oregon*);

My physician license Number is # _____ (MUST BE COMPLETED TO BE A VALID PRESCRIPTION); and I am acting within my scope of practice and authorized by law to order Infusion of the medication described above for the patient identified on this form.

Provider signature: _____ Date/Time: _____

Printed Name: _____ Phone: _____ Fax: _____

Please check the appropriate box for the patient's preferred clinic location:

Hillsboro Medical Center
 Infusion Services
 364 SE 8th Ave, Medical Plaza Suite 108B
 Hillsboro, OR 97123
 Phone number: (503) 681-4124
 Fax number: (503) 681-4120

Adventist Health Portland
 Infusion Services
 10123 SE Market St
 Portland, OR 97216
 Phone number: (503) 261-6631
 Fax number: (503) 261-6756

Mid-Columbia Medical Center
 Celilo Cancer Center
 1800 E 19th St
 The Dalles, OR 97058
 Phone number: (541) 296-7585
 Fax number: (541) 296-7610