

*Generic substitute unless checked \square

	*Patient name:			*D	OB:
	*Diagnosis and ICD-10 code:				
	Allergies:				
<u> </u>	Outpatient admit:	Series 🗆 O	ne time		
	Code status: ☐ Full code ☐ DNR ☐ Medications only ☐ Other (specify)				
	Vital signs:	□ Per protocol	☐ Other (spec	cify)	
	Access port per protocol (using sterile technique, etc.)				
	Frequency of port acces	s: One time or	der 🗖 Biweekly	√ □ Weekly □ I	Monthly
		☐ Other (speci	fy)		
	Draw blood for labs: ☐ Yes ☐ No If yes, frequency of labs:				
	Lab draws:			-	
		☐ Other (speci	fy)		
	If not drawing blood for labs, check patency of line by drawing back to get blood return and then flushing with 20 mL. NS. Follow with 5 mL 100 units/mL Heparin prior to de-accessing port.				
	If port flushes, but does not give blood return, try repositioning the patient (sitting up straighter, lying flat, a above head, etc.)				
	If port does not give blood or flush, use following protocol for Cathflo-Activase:				
	 For patient > 30 kg: Instill 2 mg/mL into the occluded lumen 				
	- For patient $<$ 30 kg: Instill volume equivalent to 100% the volume of the occluded lumen (up to 2 mg/ml				
	• May repeat the ordered dose after 120 minutes if blood still cannot be aspirated from the access device.				
	Always flush port with 5 mL 100 units/mL Heparin prior to de-accessing.				
	When finished, de-access per protocol.				
	Additional orders:				



Physician Order: Port Flush Adventist Health Tillamook 1000 Third St., Tillamook OR 97141 { Patient label }